

Other Antidepressant Dosing

Medication	Range	Starting Dose	Highest Dose	Dosing Increase	Renal Dosing	Hepatic Dosing	Cardiac
Wellbutrin SR (BID)	100-400 mg	100 mg	400 mg	100 mg	Lower initial dose or give less often; monitor closely	Lower initial dose or give less often: monitor closely	Second dose no later than 2 p.m.
Wellbutrin XL (Qam)	150-400 mg	150 mg	450 mg	150 mg	Lower initial dose or give less often; monitor closely	Lower initial dose or give less often: monitor closely	
Remeron (Qhs)	7.5-45 mg	7.5 mg	45 mg	7.5 mg	Use with caution	Use with caution; lower dose may be necessary	Monitor for sedation
Viibryd (Qam)	10-40 mg	10 mg	40 mg	10 mg	Same dosing	Same dosing for mild/mod; not studied in severe disease	
Trintellix (Qam)	5-20 mg	5 mg	20 mg	5 mg	Same dosing	Same dosing for mild/mod; not studied in severe disease	
Desyrel (Qhs)	25-200 mg	25 mg	200 mg	25 mg	Same dosing	Use with caution	Age 65+ may have adverse ↑ effects and may require lower dosing; may cause arrhythmias
Olepto (Qhs)	75-375 mg	75 mg	375 mg	75 mg	Same dosing	Use with caution	Age 65+ may have adverse ↑ effects and may require lower dosing; may cause arrhythmias

Some patients may tolerate lower doses better for all antidepressants in this chart.

Notes:



www.atomAlliance.org

This material was prepared by atom Alliance, the Quality Innovation Network-Quality Improvement Organization (QIN-QIO) coordinated by Qsource for Tennessee, Kentucky, Indiana, Mississippi and Alabama under a contract with the CENTERS for MEDICARE & MEDICAID SERVICES (CMS), a federal agency of the U.S. Department of Health and Human Services. Content does not necessarily reflect CMS policy. 18.ASG1.11.095