

# PNEUMOCOCCAL VACCINE TIMING



## Age 65 Years or Older – Everyone

- If PCV13 was given before age 65 years, no additional PCV13 is needed.

No history of pneumococcal vaccine

**PCV 13**  
PREVNAR 13®

12 month interval

**PPSV 23**  
PNEUMOVAX® 23

Received PPSV23 before age 65

1 year interval

**PCV 13**

12 month interval  
(and at least 5 years after prior dose of PPSV23)

**PPSV 23**

Received PPSV23 at age 65 or older

1 year interval

**PCV 13**

## Age 19-64 Years – Underlying Conditions

- Prior doses count towards doses recommended below and do not need to be repeated.
- If PPSV23 given previously – wait one year before giving PCV13 – when dose indicated, wait at least five years before giving a second dose of PPSV23

Smoker, long-term facility or chronic conditions:

- heart disease (excluding hypertension)
- lung disease (including asthma)
- liver disease (including cirrhosis)
- Diabetes
- alcoholism

**PPSV 23**

Immunocompromised (including HIV infection), **chronic renal failure, nephrotic syndrome or asplenia**

**PCV 13**

8 week interval

**PPSV 23**

5 year interval

**PPSV 23**

CSF leaks or cochlear implants

**PCV 13**

8 week interval

**PPSV 23**

**DO NOT ADMINISTER PCV13 AND PPSV23 AT THE SAME VISIT.**