

Frequently Asked Questions

The Facts About Fluid

“Make sure you are watching your fluids” is something dialysis patients hear all the time. But why is fluid intake so important to your health? Here are some frequently asked questions and answers about this important topic.



Why do I have to limit my daily fluid intake?

- Dialysis only removes some of the fluid
- Greater and undesired fluid gain between treatments increases risk of cardiovascular death

What are the different types of fluid?

Examples of where you find fluid:

- Water, juice, milk
- Coffee and tea
- Soups and broths
- Watermelon and other fruits
- Ice cubes, ice cream, popsicles
- Puddings, yogurt, gelatin
- Lettuce

What are potential symptoms of fluid overload?

- Shortness of breath
- Headaches
- Stomach bloating
- High blood pressure
- Heart failure
- Swelling (feet, ankles, hands)

What can happen during treatment if you remove too much fluid?

- Low blood pressure
- Cramping
- Nausea/vomiting
- Headache
- Dizziness
- Fatigue

For more information or to file a grievance, please contact:

ESRD Network 10 (IL)
911 E. 86th St., Suite 202 | Indianapolis, IN 46240
Toll Free Patient Line (800) 456-6919

ESRDNetwork10@qsource.org

ESRD Network 12 (IA, KS, MO, NE)
920 Main, Suite 801 | Kansas City, MO 64105
Toll Free Patient Line (800) 444-9965

ESRDNetwork12@qsource.org



ESRDNetwork10.org | ESRDNetwork12.org

This resource was developed while under contract with the Centers for Medicare & Medicaid Services (CMS), an agency of the U.S. Department of Health and Human Services. Contract # HHSM-500-2016-00010C and #HHSM-500-2016-00012C. The contents presented do not necessarily reflect CMS policy. 20.Q-ESRD.07.090